



# Morphine

Information for patients

UHN

## Read this medicine fact sheet to learn about:

- What this medicine is
- Why you need to take it
- What side effects to watch for

## What is the name of my medicine?

Morphine (pronounced MOR - feen)

## Other names for this medicine:

Doloral	Kadian®
M-Eslon®	M.O.S.-SR®
M.O.S.-Sulfate®	Morphine SR
Morphine-EPD	MS Contin SRT
MS Contin®	MS-IR®
Novo-Morphine SR	PMS-Morphine Sulfate SR
ratio-Morphine	ratio-Morphine SR
Sandoz-Morphine SR	Statex®
Teva-Morphine SR	

Please visit the UHN Patient Education website for more health information: [www.uhnpatienteducation.ca](http://www.uhnpatienteducation.ca)  
© 2013 University Health Network. All rights reserved.  
This information is to be used for informational purposes only and is not intended as a substitute for professional medical advice, diagnosis or treatment. Please consult your health care provider for advice about a specific medical condition. A single copy of these materials may be reprinted for non-commercial personal use only.

Author:s Adapted from Lexicomp® / Contact: Sandra Mills and Anthony Daniel  
Created: 06/2013  
Form: D-5979

## Why do I need to take this medicine?

- To ease pain.



Remember, this medicine may be addictive if you take it for a long time. You will be monitored to make sure you do not become addicted to this medicine. If you have a history of a drug or drinking problem, talk with your doctor.

This medicine may slow down your breathing too much.  
Talk with your doctor.

## How do I take this medicine?

Long-acting tablets:

- Swallow the tablets whole. Do not chew, break or crush them.
- Do not take this medicine with other strong pain medicine or if you are using a pain patch. Talk to your doctor first.
- Take it with food if it upsets your stomach.
- Drink lots of liquids with no caffeine unless your doctor tells you to drink less liquid.
- Keep a pain diary. This means writing down what your level of pain is and how much pain medicine you take.
- If you have been taking this medicine for many weeks, talk to your doctor first before you stop. You and your doctor may talk about how to slowly decrease the amount you take.
- If you have seizures, talk with your doctor.

**Tell your doctor if you have any allergies.**

## What are the side effects of this medicine?

- Feeling lightheaded, sleepy, having blurred eyesight, or a change in thinking clearly
- Dizziness
- Upset stomach or throwing up
- Hard stools or constipation. Drink more liquids, exercise or eat foods high in fibre. Talk with your doctor about taking a stool softener or laxative.

## Call your doctor right away if you notice:

- **Signs of an allergic reaction:** rash, hives, itching, red, swollen, blistered, or peeling skin with or without fever, wheezing, tightness in the chest or throat, trouble breathing or talking, or swelling of the mouth, face, lips, tongue or throat
- Poor pain control
- Side effect that is not better or you are feeling worse

**Call your doctor if you have any questions about the side effects.  
Go to your local emergency room if you think there has been an overdose.**

## What should I do if I miss a dose?

- Take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.

## **Who do I talk to if I have a problem or question?**

This pamphlet does not cover all of the side effects that may happen. Please contact a pharmacist or other health care professionals if you have any questions about this medicine or the side effects.

This site does not provide medical advice.

These sheets have been reformatted from Lexicomp<sup>®</sup>. They are provided for informational purposes only. These sheets are meant to supplement a discussion with the pharmacist.